


BENZODIAZEPINES

Benzodiazepines are a class of drugs that act on the central nervous system (CNS) to reduce activity by targeting the GABA receptor.^{1,2} Benzodiazepines are the most commonly prescribed class of drugs in the world for the treatment of insomnia and anxiety; they are also used as anticonvulsants.^{3,4} When used as prescribed, benzodiazepines are relatively safe, but have abuse potential and enhance CNS and respiratory depression when used concomitantly with opioids analgesics.⁵

Benzodiazepines are comprised of similar benzene and diazepine ring structures, making detection with immunoassay less sensitive and specific for identifying the individual drugs within the drug class. Precision Diagnostics offers our LC-MS/MS NextGen cutoffs for the individual parent drugs and metabolites listed in the chart below.

Although similar in structure, benzodiazepines exhibit different pharmacokinetic characteristics that ultimately affect the detection time in urine and overall detection in oral fluid.⁴ The lipophilic nature of benzodiazepines allows for extended detection windows in urine.⁶ Detection in oral fluid is typically less successful, due in large part to the tendency for protein binding of benzodiazepines in the blood and the glucuronidation of the associated metabolites, resulting in low concentrations in the oral fluid.⁴



† Parent drug not included in test menu

**A Precision Diagnostics trained Clinical Support Specialist can
assist with further review of your patient's results**

(800) 635-6901 Option 2

References:

1. Oshiro, C., Benzodiazepine pathway, pharmacodynamics. <https://www.pharmgkb.org/pathway/PA165111376>. Accessed May 30, 2018. ©2001-2018 PharmGKB.
2. Oshiro, C., Benzodiazepine pathway, pharmacokinetics. <https://www.pharmgkb.org/pathway/PA165111375>. Accessed May 30, 2018. ©2001-2018 PharmGKB.
3. Olsson, M., King, M., Schoenbaum, M., (2015) Benzodiazepine use in the United States. *JAMA Psychiatry*, 72(2): 136-142.
4. Allen, K. R., (2011) Screening for drugs of abuse: which matrix, oral fluid or urine? *Annals of Clinical Biochemistry*, 48: 531-541.
5. U.S. Food and Drug Administration, FDA warns about serious risks and death when combining opioid pain or cough medicines with benzodiazepines; requires its strongest warning. <https://www.fda.gov/Drugs/DrugSafety/ucm518473.htm>. Accessed May 30, 2018.
6. Nordal, K., Oiestad, E. L., Enger, A., Christophersen, A. S., Vindenes, V. (2015) Detection times of diazepam, clonazepam, and alprazolam in oral fluid collection from patients admitted to detoxification, after high and repeated drug intake. *Ther Drug Monit*, 37: 451-460.

Precision Diagnostics is a leader in clinical laboratory testing and medication adherence monitoring. Specializing in qualitative and quantitative drug testing, our innovative state-of-the art technology provides new levels of data visibility and pricing transparency.

Precision's role is to ensure each participant, from the patient to the provider and the payer, benefits from our continued commitment to the principles of value-based care and medically necessary test utilization.

Precision Diagnostics

4215 Sorrento Valley Blvd.
Suite 100
San Diego, CA 92121
(800) 635-6901
info@precisiondxlab.com
precisiondxlab.com



innovation



integrity



insights



outcomes